

## SUPPLEMENT - KETOGENIC DIET AND TREATMENTS

# Ketone bodies, glycolysis, and $K_{ATP}$ channels in the mechanism of the ketogenic diet

Gary Yellen

Department of Neurobiology, Harvard Medical School, Boston, Massachusetts, U.S.A.

### SUMMARY

The ketogenic diet (KD) has shown remarkable efficacy in the treatment of drug-resistant childhood epilepsy. Our understanding of how the KD produces its anticonvulsant and antiepileptogenic effects remains incomplete, which is perhaps not surprising for a biological manipulation as sweeping as dietary change. Several hypotheses focus on ketone bodies, fuel molecules that circulate at mil-

limolar concentrations in the blood of patients on a KD, as causative agents. Here I consider some recent evidence for one such hypothesis, involving a possible role for altered glycolysis and consequent activation of a class of potassium channels called  $K_{ATP}$  channels.

**KEY WORDS:** Ketogenic diet,  $K_{ATP}$  channels, Glycolysis, Ketone bodies.

Various mechanisms have been hypothesized for the anticonvulsant effect of the KD, differing in their focus on one or another of the many metabolic changes likely to occur in patients on the diet (Schwartzkroin, 1999; Bough & Rho, 2007). Many of these hypotheses focus on ketone bodies as crucial mediators of the beneficial effects of the KD. In a patient or animal on a KD, the liver metabolizes fats to acetoacetate, a four-carbon  $\beta$ -keto acid. Acetoacetate is enzymatically reduced to R- $\beta$ -hydroxybutyric acid, and these two so-called “ketone bodies” circulate at millimolar levels in the blood, and can serve as an energy source to all extrahepatic tissues including the brain. Acetoacetate can spontaneously decarboxylate to produce acetone, which produces the characteristic breath odor of those in ketosis. In most studies, some level of ketosis (elevation of blood ketone bodies) seems to be important for diet efficacy, though there is not necessarily a correlation between the level of ketone bodies and the degree of seizure control.

Some hypotheses for the role of ketone bodies focus on the possibility of their direct pharmacologic effects (Likhodii et al., 2003), while others focus on their role as metabolic fuels (Ma et al., 2007; Yudkoff et al., 2007). In vitro studies of ketone bodies as pharmacologic anticonvulsants have been disappointing (Thio et al., 2000), though at high concentrations, acetone and acetoacetate can have acute anticonvulsant effects in vivo (Rho et al.,

2002; Likhodii et al., 2003). An alternative role for ketone bodies—not as drugs but as metabolic fuels—is supported by early work on the metabolic effects of fasting in humans. Owen et al. (1967) showed that in fasted humans, the brain can extract ketone bodies from blood sufficient for ~65% of its energy utilization, with a corresponding reduction in the utilization of glucose. The brain’s ability to extract and use ketone bodies depends on monocarboxylate transporters in the blood–brain barrier and on specific enzymes in brain mitochondria, both of which can be induced by fasting or a ketogenic diet.

### Rapid reversal of anticonvulsant protection suggests a fast physiological mechanism

The therapeutic effects of a KD for epilepsy can be apparent quite quickly or may require days. This variation may be consistent with the need for induction of the transporters and enzymes required for ketone body utilization. On the other hand, the reversal of seizure protection when human patients break a KD can be nearly immediate—after eating a candy bar or upon IV infusion of glucose (Huttenlocher, 1976), seizure activity in patients prone to frequent seizures can recur within the hour.

This rapid reversal suggests an important role for acute physiological processes in the anticonvulsant effect of the KD. Upon glucose infusion, the levels of circulating ketone bodies decline on a fast time scale of tens of minutes (Huttenlocher, 1976; except that brain acetone requires many hours to fall; Jones, 2000). Slower mechanisms involving changes in gene and protein expression (Sullivan et al., 2004; Bough et al., 2006; Garriga-Canut et al., 2006)

Address correspondence to Gary Yellen, Ph.D., Department of Neurobiology, Harvard Medical School, Boston, MA 02115, U.S.A. E-mail: gary.yellen@hms.harvard.edu

may be important in other therapeutic effects of the KD on epileptogenesis or neuroprotection, but the anticonvulsant effect of the KD appears to involve rapidly responsive mechanisms.

### $K_{ATP}$ channels can couple metabolism and excitability

A well-known mechanism for rapid coupling between metabolism and electrical excitability is a class of potassium channels sensitive to ATP: the  $K_{ATP}$  channels. These channels play a key role in regulating insulin secretion from pancreatic beta cells. Between meals, the activity of these channels maintains the membrane potential of beta cells at a negative value; upon ingesting a carbohydrate-rich meal, the consequent elevation of intracellular ATP leads these channels to close, allowing action potential firing that promotes insulin secretion.

$K_{ATP}$  channels are also present in many central neurons, but their role is less well understood. Mice with a knock-out of the Kir6.2 pore-forming subunit of  $K_{ATP}$  channels exhibit increased sensitivity to anoxia-induced seizures (Yamada et al., 2001), implying that (at least under this condition of extreme metabolic stress)  $K_{ATP}$  channels can help to prevent excessive and harmful neuronal firing.

Although whole-brain ATP levels are elevated in animals on the KD, the oxidative metabolism of ketone bodies causes a reduction in brain glucose utilization (DeVivo et al., 1978). It has been argued that ATP production from glycolysis (the first phase of glucose utilization) plays a privileged role in controlling processes at the plasma membrane, including the regulation of  $K_{ATP}$  channels and the fueling of the ATP-driven  $Na^+$  pumps (see references in Ma et al., 2007). We therefore considered the possibility that when a brain slice metabolizes ketone bodies, the reduction in glycolytic ATP production might enable the opening of  $K_{ATP}$  channels and thus could reduce the electrical excitability of central neurons.

For our experiments, we examined the regular spontaneous firing activity of neurons in the substantia nigra pars reticulata (SNpr) (Ma et al., 2007). The GABAergic projection neurons of SNpr contain a high density of  $K_{ATP}$  channels, and the SNpr is thought to act as a “seizure gate” that regulates seizure threshold even for seizures that originate elsewhere in the brain (Iadarola & Gale, 1982; McNamara et al., 1984; Depaulis et al., 1994; Velíšková & Moshé, 2006). We used brain slices from suckling mice or rats (postnatal day 13–15) because these animals are adapted to high fat mother’s milk that approximates a KD, so the cells have all of the metabolic machinery needed to metabolize ketone bodies (Nehlig, 1999).

We found that supplying ketone bodies (acetoacetate or R- $\beta$ -hydroxybutyrate) to these brain slices, in the constant presence of glucose, produced a  $\sim 15\%$  reduction in the spontaneous firing rate, over 15–30 min of exposure. This small effect became roughly twice as big for faster firing cells, either for cells that naturally fired at a higher rate,

or for cells that were stimulated to fire by lowering extracellular  $[Ca^{2+}]$ . This “use dependence”—that is, a larger effect on faster-firing cells—is a common attribute of successful therapeutic interventions for epilepsy, as it allows for a minimal effect on normal brain activity but a strong suppression of epileptiform activity.

These effects of ketone bodies on action potential firing are eliminated in the presence of  $K_{ATP}$  blockers, or when the Kir6.2 subunit is knocked out genetically. The data support the idea that  $K_{ATP}$  channels are activated in the presence of ketone bodies, but the mechanism of this activation is not yet clear. The time course seems too slow for a direct pharmacologic effect of the ketone bodies on the channels, but it is consistent with a change in cellular metabolism.

ATP compartmentation—the notion that glycolytic ATP production is especially critical for plasma membrane processes—would explain the basic effect that ketone bodies tend to activate  $K_{ATP}$  channels, as glycolysis is reduced when ketone bodies are metabolized. It could also explain the apparent use-dependence, as increased firing allows more  $Na^+$  to enter the cell. The  $Na^+$  must be pumped out by the plasma membrane  $Na^+$  pump, thus consuming ATP in the same submembrane compartment where ATP is produced by glycolysis and sensed by  $K_{ATP}$  channels. Indeed, there is evidence from brainstem respiratory neurons that firing activity can provoke  $K_{ATP}$  channel opening by a mechanism involving the  $Na^+$  pump (Haller et al., 2001).

$K_{ATP}$  channels may thus be part of a normal negative-feedback mechanism—and perhaps a built-in seizure protection mechanism—whose set-point is determined by the level of glycolysis and can be manipulated by ketone body metabolism. Such a mechanism, if it does operate, is likely not restricted to the SNpr, as many central neurons express  $K_{ATP}$  channels. Preliminary evidence on hippocampal dentate granule neurons suggests that  $K_{ATP}$  channel open probability is increased by action potential firing, and that both induced opening and the basal open probability are augmented in the presence of ketone bodies (G. Tanner and G. Yellen, unpublished).

Under this  $K_{ATP}$ -glycolysis hypothesis, the key effect of ketone bodies is to reduce glycolytic activity. Reduced glycolysis is also an apparent trigger step for certain changes in gene expression that can retard epileptogenesis (Garriga-Canut et al., 2006). In principle, glycolysis can be reduced either by inhibiting it with a compound like 2-deoxyglucose, by lowering glucose, or by providing an oxidative substrate (like ketone bodies). The advantage of providing an oxidative substrate is that it permits a switch to an alternative fuel source without compromising the option to use glycolysis in case of metabolic stress.

It is not known exactly how glucose and ketone body concentrations interact to determine the level of glycolysis in neurons. Clinical management of blood glucose levels (as targeted directly in the low-glycemic index treatment [LGIT]; Pfeifer & Thiele, 2005) does appear to play an

important role in the efficacy of dietary treatment for epilepsy. Lowered glucose and modestly elevated ketone body concentrations (as seen in patients on the LGIT) may synergistically reduce glycolysis. Even without high ketone body concentrations, it is possible that overall flux of ketone bodies (synthesis by liver and consumption by brain) is elevated. In addition, given a diet heavier in fats than in carbohydrates, it is possible that local ketogenesis by astrocytes (Cullingford et al., 2002) could play a role in changing the mix of fuels consumed by neurons in the brain.

### The $K_{ATP}$ —glycolysis hypothesis requires further exploration

Additional studies are needed both at the cellular level, to understand the mechanism of ketone body effects on glycolysis and  $K_{ATP}$  channels, and at higher levels (brain slices, animals, and people) to discern its role in the anticonvulsant mechanism of the KD. At the cellular level, we have recently developed a novel genetically encoded fluorescent sensor of the intracellular ATP/ADP ratio (J. Berg and G. Yellen, unpublished). This tool, particularly when it is targeted to the submembrane compartment and elsewhere in neurons, should permit a more direct assessment of whether ATP compartmentation exists.

At higher levels, the most direct but also problematic approach would be to test the effect of  $K_{ATP}$  channel knockout on the efficacy of KD treatment in a mouse seizure model. The problem is that in every rodent seizure model tested, the efficacy of KD treatment has been extremely sensitive to mouse or rat strain, to age, and to the details of the diet and seizure model. The KD appears to be much less robust in treating rodent seizures than human seizures. Perhaps if it is possible to do genetic testing on a large enough cohort of human patients, then genetic contributions to efficacy—including, for instance, polymorphisms in  $K_{ATP}$  channel genes that affect the risk of diabetes, possibly due to their effect on channel open probability (Gloyn et al., 2004)—may provide additional clues to the actual mechanisms of protection in humans.

## ACKNOWLEDGMENT

I confirm that I have read the Journal's position on issues involved in ethical publication and affirm that this report is consistent with those guidelines. This work was supported by NIH/NINDS grant R01 NS055031 and a grant from the Ellison Family Foundation. I am grateful to my colleagues Jim Berg, Geoffrey Tanner, Yin P. Hung, Elizabeth Thiele, and Heidi Pfeifer for discussions.

Disclosure: The author declares no conflicts of interest.

Conflict of interest: The author has read the journal's policy on ethical publishing and agrees that this manuscript conforms to those guidelines. The author of this article discloses that there are no conflicts of interest.

## REFERENCES

- Bough KJ, Wetherington J, Hassel B, Pare JF, Gawryluk JW, Greene JG, Shaw R, Smith Y, Geiger JD, Dingledine RJ. (2006) Mitochondrial biogenesis in the anticonvulsant mechanism of the ketogenic diet. *Ann Neurol* 60:223–235.
- Bough KJ, Rho JM. (2007) Anticonvulsant mechanisms of the ketogenic diet. *Epilepsia* 48:43–58.
- Cullingford TE, Eagles DA, Sato H. (2002) The ketogenic diet upregulates expression of the gene encoding the key ketogenic enzyme mitochondrial 3-hydroxy-3-methylglutaryl-CoA synthase in rat brain. *Epilepsy Res* 49:99–107.
- Depaulis A, Vergnes M, Marescaux C. (1994) Endogenous control of epilepsy: the nigral inhibitory system. *Prog Neurobiol* 42:33–52.
- DeVivo DC, Leckie MP, Ferrendelli JS, McDougal Jr. DB. (1978) Chronic ketosis and cerebral metabolism. *Ann Neurol* 3:331–337.
- Garriga-Canut M, Schoenike B, Qazi R, Bergendahl K, Daley TJ, Pfender RM, Morrison JF, Ockuly J, Stafstrom C, Sutula T, Roopra A. (2006) 2-Deoxy-D-glucose reduces epilepsy progression by NRSF-CtBP dependent metabolic regulation of chromatin structure. *Nat Neurosci* 9:1382–1387.
- Gloyn AL, Pearson ER, Antcliff JF, Proks P, Bruining GJ, Slingerland AS, Howard N, Srinivasan S, Silva JM, Molnes J, Edghill EL, Frayling TM, Temple IK, Mackay D, Shield JP, Sumnik Z, van Rhijn A, Wales JK, Clark P, Gorman S, Aisenberg J, Ellard S, Njolstad PR, Ashcroft FM, Hattersley AT. (2004) Activating mutations in the ATP-sensitive potassium channel subunit Kir6.2 gene are associated with permanent neonatal diabetes. *New Engl J Med* 350:1838–1849.
- Haller M, Mironov SL, Karschin A, Richter DW. (2001) Dynamic activation of  $K_{ATP}$  channels in rhythmically active neurons. *J Physiol Lond* 537:69–81.
- Huttenlocher PR. (1976) Ketonemia and seizures: metabolic and anticonvulsant effects of two ketogenic diets in childhood epilepsy. *Pediatr Res* 10:536–540.
- Iadarola MJ, Gale K. (1982) Substantia nigra: site of anticonvulsant activity mediated by gamma-aminobutyric acid. *Science* 218:1237–1240.
- Jones AW. (2000) Elimination half-life of acetone in humans: case reports and review of the literature. *J Anal Toxicol* 24:8–10.
- Likhodii SS, Serbanescu I, Cortez MA, Murphy P, Snead OC, Burnham WM. (2003) Anticonvulsant properties of acetone, a brain ketone elevated by the ketogenic diet. *Ann Neurol* 54:219–226.
- Ma W, Berg J, Yellen G. (2007) Ketogenic diet metabolites reduce firing in central neurons by opening  $K_{ATP}$  channels. *J Neurosci* 27:3618–3625.
- McNamara JO, Galloway MT, Rigsbee LC, Shin C. (1984) Evidence implicating substantia nigra in regulation of kindled seizure threshold. *J Neurosci* 4:2410–2417.
- Nehlig A. (1999) Age-dependent pathways of brain energy metabolism: the suckling rat, a natural model of the ketogenic diet. *Epilepsy Res* 37:211–221.
- Owen OE, Morgan AP, Kemp HG, Sullivan JM, Herrera MG, Cahill Jr. GF. (1967) Brain metabolism during fasting. *J Clin Invest* 46:1589–1595.
- Pfeifer HH, Thiele EA. (2005) Low-glycemic-index treatment: a liberalized ketogenic diet for treatment of intractable epilepsy. *Neurology* 65:1810–1812.
- Rho JM, Anderson GD, Donevan SD, White HS. (2002) Acetoacetate, acetone, and dibenzylamine (a contaminant in L-(+)- $\beta$ -hydroxybutyrate) exhibit direct anticonvulsant actions in vivo. *Epilepsia* 43:358–361.
- Schwartzkroin PA. (1999) Mechanisms underlying the anti-epileptic efficacy of the ketogenic diet. *Epilepsy Res* 37:171–180.
- Sullivan PG, Rippey NA, Dorenbos K, Concepcion RC, Agarwal AK, Rho JM. (2004) The ketogenic diet increases mitochondrial uncoupling protein levels and activity. *Ann Neurol* 55:576–580.
- Thio LL, Wong M, Yamada KA. (2000) Ketone bodies do not directly alter excitatory or inhibitory hippocampal synaptic transmission. *Neurology* 54:325–331.
- Velísková J, Moshé SL. (2006) Update on the role of substantia nigra pars reticulata in the regulation of seizures. *Epilepsy Curr* 6:83–87.
- Yamada K, Ji JJ, Yuan H, Miki T, Sato S, Horimoto N, Shimizu T, Seino S, Inagaki N. (2001) Protective role of ATP-sensitive potassium channels in hypoxia-induced generalized seizure. *Science* 292:1543–1546.
- Yudkoff M, Daikhin Y, Melo TM, Nissim I, Sonnewald U, Nissim I. (2007) The ketogenic diet and brain metabolism of amino acids: relationship to the anticonvulsant effect. *Annu Rev Nutr* 27:415–430.